

Deuteron Anapole Moment with Heavy Mesons

C.-P. Liu^{1,*}, C. H. Hyun^{2,†} and B. Desplanques^{3‡}

¹*TRIUMF, 4004 Wesbrook Mall, Vancouver, British Columbia, Canada V6T 2A3*

²*Institute of Basic Science, Sungkyunkwan University, Suwon 440-746, Korea*

³*Laboratoire de Physique Subatomique et de Cosmologie
(UMR CNRS/IN2P3-UJF-INPG), F-38026 Grenoble Cedex, France*

(Dated: 8th February 2008)

Abstract

Parity-nonconserving two-body currents due to vector meson exchange are considered with the aim to determine the related contributions to the anapole moment. A particular attention is given to the requirement of current conservation which is essential for a reliable estimate of this quantity. An application is made for the deuteron case.

*Electronic address: cpliu@triumf.ca

†Electronic address: hch@meson.skku.ac.kr

‡Electronic address: desplanq@lpsc.in2p3.fr

I. INTRODUCTION

First introduced by Zel'dovich [1], the anapole moment (AM) of a quantum system is a quantity that involves both the electromagnetic interaction and parity nonconservation (PNC). Ignored for a long time, it is not before studies by Flambaum and Khriplovich [2] that the concept acquires practical interest. In their work, these authors especially emphasize that the AM would grow with the size of the nucleus, making heavy nuclei natural candidates for an observation. Implying the hyperfine structure, a first measurement was performed a few years ago in the ^{133}Cs nucleus [3].

The deuteron AM has also received some attention recently [4, 5, 6, 7, 8]. Though its interest is largely academic (an experiment is not feasible in a near future), it offers the advantage of a laboratory where methods and ingredients relative to an estimate can be studied in details. These studies have been concerned with the pion-exchange component of the PNC nucleon-nucleon (NN) force. Calculations were based on assuming an effective-field-theory description [4, 5], zero-range NN strong forces [7], or more realistic NN strong forces [8]. The use of an alternative field-theory description was also proposed [6]. The next step concerns the extension of these results to include the component of the PNC force due to vector-meson exchange that could contribute as much as, if not more than the pion exchange.

Determining the AM supposes to calculate the effective current that couples to the photon. As it involves parity nonconservation, the current has necessarily an axial character, making the requirement of current conservation non-trivial. Individual contributions are proportional to the weak coupling, G_F , while fulfilling the above property implies that the effective current contains the factor $G_F q^2$, which vanishes in the limit of a zero momentum transfer. Getting this result demands particular care with contributions ensuring gauge invariance. When dealing with vector mesons, this task becomes essential. In particular, it has to be done consistently with the PNC interaction model that is employed in calculations. Some contributions were given in Ref. [9]. In the present work, we intend to complete this work with the double aim to satisfy gauge invariance and consistency with the PNC interaction model; DDH potential, given by Desplanques, Donoghue, and Holstein, is our case [10]. Some estimates of the vector-meson-exchange contributions to the anapole moment will be presented in the deuteron case.

The plan of the paper is as follows. In Section II, we first present the various ingredients pertinent to the interaction: parity-conserving (PC), parity-nonconserving and electromagnetic (EM) ones. We subsequently provide the expressions for the PNC two-body currents at the lowest $1/m_N$ order and show how they allow one to fulfill current conservation. Section III is devoted to applications

involving the deuteron. This includes the deuteron description, especially the determination of the PNC components; the expression of the anapole matrix elements from both the one- and two-body currents; and a numerical estimate in terms of the PNC meson-nucleon coupling constants. A discussion of the results is given in Section IV. This is completed by Appendix A that contains expressions of the two-body currents in configuration space.

II. PNC NN INTERACTION, CURRENTS, AND CURRENT CONSERVATION

The anapole moment is a special electromagnetic property of a system in which parity conservation is violated; therefore, our first step in calculation is formulating the EM current operators. Throughout this work, we assume for a nuclear system the validity of non-relativistic (NR) limit and keep only terms of leading order.

The one-body term ($\rho^{(1)}, \mathbf{j}^{(1)}$), that is, contributions from each individual nucleon (for deuteron, there are two), include the charge density:

$$\rho^{(1)} = e \sum_{i=1}^2 \frac{1 + \tau_i^z}{2} (2\pi)^3 \delta^{(3)}(\mathbf{k} + \mathbf{p}'_i - \mathbf{p}_i), \quad (1)$$

and the 3-current densities from spin and motion of nucleons:

$$\mathbf{j}_{spin} = e \sum_{i=1}^2 \frac{\mu_N^i}{2m_N} i\boldsymbol{\sigma}_i \times (\mathbf{p}'_i - \mathbf{p}_i) (2\pi)^3 \delta^{(3)}(\mathbf{k} + \mathbf{p}'_i - \mathbf{p}_i), \quad (2)$$

$$\mathbf{j}_{conv} = e \sum_{i=1}^2 \frac{1 + \tau_i^z}{4m_N} (\mathbf{p}'_i + \mathbf{p}_i) (2\pi)^3 \delta^{(3)}(\mathbf{k} + \mathbf{p}'_i - \mathbf{p}_i), \quad (3)$$

in momentum space, where μ_N^i is defined as

$$\mu_N^i \equiv \frac{1}{2} (\mu_S + \tau_i^z \mu_V)$$

with $\mu_S = 0.88$ and $\mu_V = 4.71$; and $\mathbf{k}$, $\mathbf{p}'_i$, and $\mathbf{p}_i$ denote the 3-momentum of outgoing photon, outgoing i th nucleon, and incoming i th nucleon, respectively.

As the canonical picture of nucleon-nucleon (NN) interaction is realized through the exchange of mesons, the nuclear EM currents should also contain these exchange effects, which are two-body in character. Since we are still lacking of a fundamental understanding about the NN interaction, the exact exchange currents (ECs) are also unknown. However, in order to reduce theoretical uncertainties in the studies of nuclear EM processes, it is important, given a chosen model for the NN interaction, to construct these two-body currents which are constrained by both current conservation and phenomenology.

For the PC NN interaction, many high-quality model potentials exist, and our choice for calculation in this work is the Argonne v_{18} potential (Av_{18}) [11]. Although it gives good fits to the scattering data and deuteron properties, it is not straightforward to construct the corresponding ECs because the connection with the meson exchange picture is not clear for some parts in this potential. One traditional way to construct the ECs is implementing the NR minimal coupling (MC) to the potential, i.e.,

$$\mathbf{p} \rightarrow \mathbf{p} - \frac{e}{2}(1 + \tau^z)\mathbf{A}; \quad H \rightarrow H + \frac{e}{2}(1 + \tau^z)A^0,$$

then identifying the EM currents from the interaction Hamiltonian density, $ej_\mu A^\mu$. However, this procedure only constrains the longitudinal components, while giving no information about the transverse components which are conserved by themselves. Some uncertainty about these last terms comes from potentials involving quadratic velocity-dependent components as discussed in Ref. [12]. Moreover, the derivation of exchange currents for a model like Av_{18} , employed in Ref. [13] for instance, requires further elaboration. This model contains a Gaussian type component while the above prescription is usually applied to Yukawa potentials. Therefore, we leave this as an open question for future work and follow the treatment of Ref. [8] to examine: (1) to what degree current conservation is broken by the omission of PC ECs, and (2) how much the inclusion of PC ECs due to the one pion exchange, which gives the long-range part in Av_{18} , could restore the conservation. By this exercise, one can get some qualitative handle on this problem. The detail is to be discussed in Section IV.

For the PNC NN interaction, our choice is the potential suggested by Desplanques, Donoghue, and Holstein (DDH) [10]. Since this potential, based on the one-boson exchange scheme involving π , ρ , and ω mesons, has a close tie with the exchange picture, a more field-theoretical formalism, the so-called S -matrix approach [9, 14, 15, 16], is used to construct all the corresponding ECs. As we will show later, some transverse components arise naturally in this derivation. For clarity, we divide the following discussion into three parts: first, the model Lagrangian, consistent with the DDH scheme, is constructed; second, the PNC ECs are derived; and finally, we show how these exchange currents allow one to fulfill current conservation, given the DDH potential.

A. Model Lagrangian

The total Lagrangian density we consider is expressed as

$$\mathcal{L} = \mathcal{L}_0 + \mathcal{L}_{PC} + \mathcal{L}_{PNC} + \mathcal{L}_{EM} + \delta\mathcal{L}, \quad (4)$$

where $\delta\mathcal{L}$ contains all the terms not relevant for this discussion. The free Lagrangian densities for nucleon, N ; pion, rho, and omega mesons, π , ρ^μ , and ω^μ , are

$$\begin{aligned} \mathcal{L}_0 = & \bar{N}'(i\not{\partial} - m_N)N + \frac{1}{2}(\partial_\mu\pi) \cdot (\partial^\mu\pi) - \frac{1}{2}m_\pi^2\pi^2 - \frac{1}{4}\mathbf{F}_{\mu\nu}^{(\rho)} \cdot \mathbf{F}^{(\rho)\mu\nu} + \frac{1}{2}m_\rho^2\rho_\mu \cdot \rho^\mu - \frac{1}{2\xi}(\partial_\mu\rho^\mu) \cdot (\partial_\nu\rho^\nu) \\ & - \frac{1}{4}F_{\mu\nu}^{(\omega)}F^{(\omega)\mu\nu} + \frac{1}{2}m_\omega^2\omega_\mu\omega^\mu - \frac{1}{2\xi}(\partial_\mu\omega^\mu)(\partial_\nu\omega^\nu), \end{aligned} \quad (5)$$

where $\mathbf{F}_{\mu\nu}^{(\rho)}$ and $F_{\mu\nu}^{(\omega)}$ are the field tensors of the vector mesons, and we keep the R_ξ gauge-fixing terms of vector mesons explicit for the moment. The PC and PNC meson-nucleon interaction Lagrangian densities are

$$\begin{aligned} \mathcal{L}_{PC} = & ig_{\pi NN}\bar{N}'\gamma_5\boldsymbol{\tau} \cdot \boldsymbol{\pi}N - g_{\rho NN}\bar{N}'(\gamma_\mu - i\frac{\chi_V}{2m_N}\sigma_{\mu\nu}q^\nu)\boldsymbol{\tau} \cdot \boldsymbol{\rho}^\mu N \\ & - g_{\omega NN}\bar{N}'(\gamma_\mu - i\frac{\chi_S}{2m_N}\sigma_{\mu\nu}q^\nu)\omega^\mu N, \end{aligned} \quad (6)$$

$$\begin{aligned} \mathcal{L}_{PNC} = & -\frac{h_\pi^1}{\sqrt{2}}\bar{N}'(\boldsymbol{\tau} \times \boldsymbol{\pi})^z N + \bar{N}'[h_\rho^0\boldsymbol{\tau} \cdot \boldsymbol{\rho}^\mu + h_\rho^1\rho^{z\mu} + \frac{h_\rho^2}{2\sqrt{6}}(3\tau^z\rho^{z\mu} - \boldsymbol{\tau} \cdot \boldsymbol{\rho}^\mu)]\gamma_\mu\gamma_5 N \\ & + \bar{N}'(h_\omega^0\omega^\mu + h_\omega^1\tau^z\omega^\mu)\gamma_\mu\gamma_5 N, \end{aligned} \quad (7)$$

where q^μ is the 4-momentum carried by the outgoing boson; the strong couplings, g_{XNN} , as well as the weak couplings, $h_X^{(i)}$, are of DDH's definition (except their PNC pion coupling, f_π , is renamed as h_π^1 here); and the anomalous strong isoscalar and isovector magnetic moments of nucleon, χ_S and χ_V , are assumed to be the same as the EM values, -0.12 and 3.70, by vector meson dominance.

Now we apply the covariant MC:

$$p_\mu \rightarrow p_\mu - \frac{e}{2}(1 + \tau^z)A_\mu,$$

to the above Lagrangian densities to obtain the EM interactions. For our purpose, only terms of first-order in e are included in $\mathcal{L}_{EM}$. From $\mathcal{L}_0$, we get

$$\mathcal{L}_{EM}^{(NN\gamma)} = -e\bar{N}'[(F_1^{(S)}(Q^2)\frac{1}{2} + F_1^{(V)}(Q^2)\frac{\tau^z}{2})\gamma_\mu - i\frac{1}{2m_N}(F_2^{(S)}(Q^2)\frac{1}{2} + F_2^{(V)}(Q^2)\frac{\tau^z}{2})\sigma_{\mu\nu}q^\nu]NA^\mu, \quad (8)$$




Figure 1: The vertex factor for vector-meson-photon coupling, where α , β , and γ are the Lorentz indices; i and j are isospin indices.

$$\mathcal{L}_{EM}^{(\pi\pi\gamma)} = -e(\boldsymbol{\pi} \times \partial_\mu \boldsymbol{\pi})^z A^\mu, \quad (9)$$

$$\mathcal{L}_{EM}^{(\rho\rho\gamma)} = -e(\boldsymbol{\rho}^\nu \times \mathbf{F}_{\nu\mu}^{(\rho)})^z A^\mu - \frac{1}{\xi} e(\boldsymbol{\rho}_\mu \times \partial_\nu \boldsymbol{\rho}^\nu)^z A^\mu. \quad (10)$$

Note that in order to account for the nucleon structure, nucleon EM form factors, $F_{1,2}^{(S,V)}$ (superscript "S" for isoscalar and "V" for isovector; subscript "1" for Dirac and "2" for Pauli) have to be added. At $Q^2 = -q^2 = 0$, $F_1^{(S)}(0) = F_1^{(V)}(0) = 1$, $F_2^{(S)}(0) = -0.12$, and $F_2^{(V)}(0) = 3.70$. In principle one should also take into account the meson structures; however, they are still poorly constrained so we simply assume them elementary.

Due to the momentum-dependent coupling of ρ meson to the nucleon anomalous magnetic moment, a Kroll-Ruderman type contact interaction [17], reading as

$$\mathcal{L}_{EM}^{(NN\rho\gamma)} = -e \frac{g_{\rho NN} \chi_V}{2m_N} \bar{N}' \sigma_{\mu\nu} (\boldsymbol{\tau} \times \boldsymbol{\rho}^\nu)^z N A^\mu, \quad (11)$$

also arises. This will lead to a seagull current which is important for current conservation, but was ignored in Ref. [9].

It is worthwhile to point out that the EM interactions obtained above depend on the model Lagrangians we start out. For example, compared with the result of QHD II [18], a gauge theory approach which first models pion from a global $SU(2)$ symmetry group and then adds ρ meson and photon by a local $SU(2) \otimes U(1)$ symmetry group (both steps involve Higgs mechanism to generate meson masses), one observes a larger $\rho\rho\gamma$ interaction in QHD II by the amount of

$$\Delta \mathcal{L}_{EM}^{(\rho\rho\gamma)} = \frac{1}{2} e (\boldsymbol{\rho}_\mu \times \boldsymbol{\rho}_\nu)^z F^{(\gamma)\mu\nu},$$

which modifies the $\rho\rho\gamma$ vertex in an interesting way as we are going to explain.

By fixing the gauge parameter $\xi = 1$, i.e., 't Hooft-Feynman gauge, we obtained the same $\mathcal{L}_{EM}^{(\rho\rho\gamma)}$ as in Ref. [19], and this gives a vertex factor (see Fig. 1):

$$\epsilon_{3ij} [(q_1 - q_2)^\mu g^{\alpha\beta} + k^\alpha g^{\beta\mu} - k^\beta g^{\mu\alpha}], \quad (12)$$

with $k + q_1 + q_2 = 0$. The latter two terms are combined to give amplitudes conserved by themselves when dotted by k_μ , which means they are purely transverse (actually, they correspond to magnetic dipole couplings which explain the transversality). By adding $\Delta\mathcal{L}_{EM}^{(\rho\rho\gamma)}$, it simply doubles the self-conserved terms, so that we have

$$\epsilon_{3ij}[(q_1 - q_2)^\mu g^{\alpha\beta} + 2k^\alpha g^{\beta\mu} - 2k^\beta g^{\mu\alpha}], \quad (13)$$

for the vertex. As ρ meson is a spin-one particle, it has charge (c), magnetic dipole (μ), and charge quadrupole (Q) couplings to the EM field. When assuming it is an elementary particle, $c = e$, $\mu = e/m_\rho$, and $Q = -e/m_\rho^2$, the vertex factor appears to be Eq. (13) [20, 21], so the MC result under-predicts the ρ meson magnetic moment by two. This factor of two difference in self-conserved terms between MC and chiral Lagrangian approaches has been pointed out in Refs. [16, 22]. It is also obtained from a quark model calculation of the charged ρ meson magnetic moment (in the limit where the ρ mass is taken as twice the constituent quark one). Although the factor two originates from different models, in order to have a closer contact with phenomenology, we make this particular modification to the $\rho\rho\gamma$ vertex.

The modification mentioned above is just an example of model-dependences in constructing ECs. Since these purely transverse terms, often called non-Born (NB) terms, could not be constrained by current conservation, it is not easy to set up criteria *a priori* to judge which should be included or not, unless compared with experiments¹. For our calculation, we choose to include $\rho\pi\gamma$ and $\omega\pi\gamma$ interactions

$$\mathcal{L}_{EM}^{(\rho\pi\gamma)} = e \frac{g_{\rho\pi\gamma}}{2m_\rho} \epsilon_{\alpha\beta\gamma\delta} F^{(\gamma)\alpha\beta} (\boldsymbol{\rho}^\gamma \cdot \partial^\delta \boldsymbol{\pi}), \quad (14)$$

$$\mathcal{L}_{EM}^{(\omega\pi\gamma)} = e \frac{g_{\omega\pi\gamma}}{2m_\omega} \epsilon_{\alpha\beta\gamma\delta} F^{(\gamma)\alpha\beta} (\boldsymbol{\omega}^\gamma \partial^\delta \pi^z), \quad (15)$$

where the total anti-symmetric tensor is defined as $\epsilon_{0123} = -1$ [15, 24], because these coupling constants could be determined from the decay data (except for signs); and we ignore all the nucleon isobaric excitations for they are not the main theoretical emphasis of this work. The present work could be easily extended to them if necessary.

To sum up, the total EM Lagrangian density we consider is

$$\mathcal{L}_{EM} = L_{EM}^{(NN\gamma)} + L_{EM}^{(\pi\pi\gamma)} + \left\{ L_{EM}^{(\rho\rho\gamma)} + \frac{1}{2} e (\boldsymbol{\rho}_\mu \times \boldsymbol{\rho}_\nu)^z F^{(\gamma)\mu\nu} \right\} + \mathcal{L}_{EM}^{(\rho\pi\gamma)} + \mathcal{L}_{EM}^{(\omega\pi\gamma)}. \quad (16)$$

¹ There are a lot of discussions in literature addressing this issue. We refer readers to more extensive reviews such as Refs. [12, 15, 16, 23] (and references therein) for details.




Figure 2: Classification of meson exchange currents: (a) norm-recoil, (b) pair, (c) mesonic, (d) seagull, (e) isobaric, and (f) non-Born mesonic, where N and N^* denote nucleon and nucleon excited state; M and M' denote mesons.

B. PNC Meson Exchange Currents

Diagrammatically, the ECs could be classified, according to Fig. 2, as: (a) norm-recoil, (b) pair, (c) mesonic, (d) seagull, (e) isobaric, and (f) NB mesonic types. The division of norm-recoil and pair terms simply comes from the separation of positive- and negative-energy components in the covariant nucleon propagator. Sometimes confusion arises when it comes to pair and seagull diagram. If the PC πNN coupling is formulated as pseudo-vector, the seagull term is $O(1/m_N)$, while the pair term is higher-order in $1/m_N$. On the other hand, if the pseudo-scalar coupling is adopted as we do here, there is no seagull term, however, at the leading order, the pair term looks exactly as the seagull term in the pseudo-vector scheme. Therefore, as far as the NR approximation is valid, these two formalisms are equivalent [12, 23]. For the case of ρ meson, both pair and seagull diagrams have $O(1/m_N)$ contributions. As for the NB contributions from (e) and (f), we only consider the latter as explained above.

Before one applies Feynman rules to evaluate these diagrams and extract the corresponding ECs, the gauge parameter has to be fixed. Though the physical results should be gauge-independent, a proper choice may greatly simplify the calculation. Here, we adopt the 't Hooft-Feynman gauge,

$\xi = 1$, for the following reasons. First, the propagator is simpler

$$\begin{aligned}\langle A_\mu A_\nu \rangle &= \left(g_{\mu\nu} - (1 - \xi) \frac{q_\mu q_\nu}{q^2 - \xi m^2 + i\epsilon} \right) \frac{-i}{q^2 - m^2 + i\epsilon} \\ &= \frac{-i g_{\mu\nu}}{q^2 - m^2 + i\epsilon} \quad (\text{for } \xi = 1).\end{aligned}$$

Second, the PNC potential, constructed from NR reduction of the one-boson exchange diagrams, corresponds to the form given by DDH. The last and most important of all, in combination with the previous statement, the contribution from the norm-recoil diagram represents how the one-body EM matrix element is modified by the presence of this NR potential [16]. Therefore, it should not be double-counted if one has already taken care of this by using the perturbed wave function, the route we will follow.

In momentum space, to the order of $1/m_N$, we have the following results for the pair and ρ -seagull (KR) 3-currents

$$\mathbf{j}_{pair}^\pi = \frac{-eg_{\pi NN} h_\pi^1}{2\sqrt{2}m_N} (\boldsymbol{\tau}_1 \cdot \boldsymbol{\tau}_2 - \tau_1^z \tau_2^z) [\boldsymbol{\sigma}_1] \frac{(2\pi)^3 \delta^{(3)}(\dots)}{\mathbf{q}_2^2 + m_\pi^2} + (1 \leftrightarrow 2), \quad (17)$$

$$\begin{aligned}\mathbf{j}_{pair+KR}^\rho &= \frac{-eg_{\rho NN}}{2m_N} \left\{ \left(h_\rho^0 (\boldsymbol{\tau}_1 \cdot \boldsymbol{\tau}_2 + \tau_2^z) + \frac{h_\rho^2}{2\sqrt{6}} (3\tau_1^z \tau_2^z - \boldsymbol{\tau}_1 \cdot \boldsymbol{\tau}_2 + 2\tau_2^z) \right) [\boldsymbol{\sigma}_1 - \boldsymbol{\sigma}_2] \right. \\ &\quad \left. + h_\rho^1 (1 + \tau_1^z) [\tau_2^z \boldsymbol{\sigma}_1 - \tau_1^z \boldsymbol{\sigma}_2] + (1 + \chi_V) \left(h_\rho^0 - \frac{h_\rho^2}{2\sqrt{6}} \right) (\boldsymbol{\tau}_1 \times \boldsymbol{\tau}_2)^z [\boldsymbol{\sigma}_1 \times \boldsymbol{\sigma}_2] \right\} \\ &\quad \times \frac{(2\pi)^3 \delta^{(3)}(\dots)}{\mathbf{q}_2^2 + m_\rho^2} + (1 \leftrightarrow 2), \quad (18)\end{aligned}$$

$$\mathbf{j}_{pair}^\omega = \frac{-eg_{\omega NN}}{2M_N} (1 + \tau_1^z) (h_\omega^0 [\boldsymbol{\sigma}_1 - \boldsymbol{\sigma}_2] + h_\omega^1 [\tau_1^z \boldsymbol{\sigma}_1 - \tau_2^z \boldsymbol{\sigma}_2]) \frac{(2\pi)^3 \delta^{(3)}(\dots)}{\mathbf{q}_2^2 + m_\omega^2} + (1 \leftrightarrow 2), \quad (19)$$

where the ρ -seagull 3-current corresponds to the term involving χ_V ; $\mathbf{q}_{1,2} = \mathbf{p}'_{1,2} - \mathbf{p}_{1,2}$; and the δ function imposes the total 3-momentum conservation, $\mathbf{k} + \mathbf{q}_1 + \mathbf{q}_2 = 0$. All the pair charges are of higher order in $1/m_N$ compared with the nucleon charge, which is $O(1)$, so they are neglected.

For the mesonic 3-currents, we obtain

$$\mathbf{j}_{mesonic}^\pi = \frac{-eg_{\pi NN} h_\pi^1}{2\sqrt{2}m_N} (\boldsymbol{\tau}_1 \cdot \boldsymbol{\tau}_2 - \tau_1^z \tau_2^z) [\mathbf{q}_2 - \mathbf{q}_1] \boldsymbol{\sigma}_1 \cdot \mathbf{q}_1 \frac{(2\pi)^3 \delta^{(3)}(\dots)}{(\mathbf{q}_1^2 + m_\pi^2)(\mathbf{q}_2^2 + m_\pi^2)} + (1 \leftrightarrow 2), \quad (20)$$

$$\begin{aligned}\mathbf{j}_{mesonic}^\rho &= \frac{-eg_{\rho NN}}{2m_N} \left(h_\rho^0 - \frac{h_\rho^2}{2\sqrt{6}} \right) i(\boldsymbol{\tau}_1 \times \boldsymbol{\tau}_2)^z \left\{ [\mathbf{q}_1 - \mathbf{q}_2] \boldsymbol{\sigma}_2 \cdot ((\mathbf{p}'_2 + \mathbf{p}_2) - (\mathbf{p}'_1 + \mathbf{p}_1) - i(1 + \chi_V) \boldsymbol{\sigma}_1 \times \mathbf{q}_1) \right. \\ &\quad \left. + 2[(\mathbf{p}'_1 + \mathbf{p}_1) + i(1 + \chi_V) \boldsymbol{\sigma}_1 \times \mathbf{q}_1] \boldsymbol{\sigma}_2 \cdot \mathbf{k} + 2[\boldsymbol{\sigma}_2] \left(2m_N k_0 - \mathbf{k} \cdot ((\mathbf{p}'_1 + \mathbf{p}_1) + i(1 + \chi_V) \boldsymbol{\sigma}_1 \times \mathbf{q}_1) \right) \right\} \\ &\quad \times \frac{(2\pi)^3 \delta^{(3)}(\dots)}{(\mathbf{q}_1^2 + m_\rho^2)(\mathbf{q}_2^2 + m_\rho^2)} + (1 \leftrightarrow 2), \quad (21)\end{aligned}$$

$$\mathbf{j}_{mesonic}^{\rho\pi} = \frac{eg_{\rho NN}g_{\rho\pi\gamma}h_{\pi}^1}{\sqrt{2}m_{\rho}}(\boldsymbol{\tau}_1 \times \boldsymbol{\tau}_2)^z[\mathbf{q}_1 \times \mathbf{q}_2] \frac{(2\pi)^3\delta^{(3)}(\dots)}{(\mathbf{q}_1^2 + m_{\rho}^2)(\mathbf{q}_2^2 + m_{\pi}^2)} + (1 \leftrightarrow 2), \quad (22)$$

$$\mathbf{j}_{mesonic}^{\omega\pi} \approx 0, \quad (23)$$

where $k_0 = E_1 + E_2 - E'_1 - E'_2 = E_i - E_f$. Specially note that there is a contribution to the charge density at the same order as the nucleon charge:

$$\rho_{mesonic}^{\rho} = -2eg_{\rho NN} \left(h_{\rho}^0 - \frac{h_{\rho}^2}{2\sqrt{6}} \right) i(\boldsymbol{\tau}_1 \times \boldsymbol{\tau}_2)^z \boldsymbol{\sigma}_2 \cdot \mathbf{k} \frac{(2\pi)^3\delta^{(3)}(\dots)}{(\mathbf{q}_1^2 + m_{\rho}^2)(\mathbf{q}_2^2 + m_{\rho}^2)} + (1 \leftrightarrow 2). \quad (24)$$

C. PNC NN Interaction and Current Conservation

By considering the one-boson exchange diagrams where one of the meson-nucleon couplings is PC and the other PNC, the DDH potential in momentum space could be expressed as

$$V_{PNC}^{\pi} = \frac{g_{\pi NN}h_{\pi}^1}{2\sqrt{2}m_N} i(\boldsymbol{\tau}_1 \times \boldsymbol{\tau}_2)^z (\boldsymbol{\sigma}_1 + \boldsymbol{\sigma}_2) \cdot \mathbf{u}_{\pi}, \quad (25)$$

$$V_{PNC}^{\rho} = \frac{-g_{\rho NN}}{m_N} \left[\left(h_{\rho}^0 \boldsymbol{\tau}_1 \cdot \boldsymbol{\tau}_2 + \frac{h_{\rho}^1}{2}(\tau_1^z + \tau_2^z) + \frac{h_{\rho}^2}{2\sqrt{6}}(3\tau_1^z\tau_2^z - \boldsymbol{\tau}_1 \cdot \boldsymbol{\tau}_2) \right) \right. \\ \left. \times (i(1 + \chi_V)(\boldsymbol{\sigma}_1 \times \boldsymbol{\sigma}_2) \cdot \mathbf{u}_{\rho} + (\boldsymbol{\sigma}_1 - \boldsymbol{\sigma}_2) \cdot \mathbf{v}_{\rho}) - \frac{h_{\rho}^1}{2}(\tau_1^z - \tau_2^z)(\boldsymbol{\sigma}_1 + \boldsymbol{\sigma}_2) \cdot \mathbf{v}_{\rho} \right], \quad (26)$$

$$V_{PNC}^{\omega} = \frac{-g_{\omega NN}}{m_N} \left[\left(h_{\omega}^0 + \frac{h_{\omega}^1}{2}(\tau_1^z + \tau_2^z) \right) (i(1 + \chi_S)(\boldsymbol{\sigma}_1 \times \boldsymbol{\sigma}_2) \cdot \mathbf{u}_{\omega} + (\boldsymbol{\sigma}_1 - \boldsymbol{\sigma}_2) \cdot \mathbf{v}_{\omega}) \right. \\ \left. + \frac{h_{\omega}^1}{2}(\tau_1^z - \tau_2^z)(\boldsymbol{\sigma}_1 + \boldsymbol{\sigma}_2) \cdot \mathbf{v}_{\omega} \right], \quad (27)$$

where

$$\mathbf{u}_X = \frac{\mathbf{q}}{q^2 + m_X^2}; \quad \mathbf{v}_X = \frac{(\mathbf{p}'_1 + \mathbf{p}_1) - (\mathbf{p}'_2 + \mathbf{p}_2)}{2(q^2 + m_X^2)},$$

with $\mathbf{q}$ denotes the meson 3-momentum.

To prove the conservation of these PNC currents at the operator level, we showed explicitly the following matrix element identities (with bra $\langle \mathbf{p}'_1, \mathbf{p}'_2 |$ and ket $| \mathbf{p}_1, \mathbf{p}_2 \rangle$):

$$\langle [\rho^{(1)}, V_{PNC}^{\pi}] \rangle = \mathbf{k} \cdot \langle \mathbf{j}_{pair}^{\pi} + \mathbf{j}_{mesonic}^{\pi} \rangle, \quad (28)$$

$$\langle [\rho^{(1)}, V_{PNC}^{\rho}] \rangle = \mathbf{k} \cdot \langle \mathbf{j}_{pair+KR}^{\rho} + \mathbf{j}_{mesonic}^{\rho(I)} \rangle, \quad (29)$$

$$\langle [\rho^{(1)}, V_{PNC}^{\omega}] \rangle = \mathbf{k} \cdot \langle \mathbf{j}_{pair}^{\omega} \rangle, \quad (30)$$

$$\langle [\rho_{mesonic}^{\rho}, H] \rangle = \mathbf{k} \cdot \langle \mathbf{j}_{mesonic}^{\rho(II)} \rangle, \quad (31)$$

$$0 = \mathbf{k} \cdot \langle \mathbf{j}_{mesonic}^{\rho\pi} + \mathbf{j}_{mesonic}^{\omega\pi} \rangle, \quad (32)$$

where H is the total Hamiltonian, which is the sum of kinetic energy (T) and both PC and PNC potentials (V_{PC} and V_{PNC}); and the ρ mesonic current, Eq. (21), is separated into two parts: (I) is proportional to the vector $(\mathbf{q}_1 - \mathbf{q}_2)$, and (II) contains the rest. The continuity equality of Eq. (31) indicates that $(\rho_{mesonic}^\rho, \mathbf{j}_{mesonic}^{\rho(II)})$ forms a conserved current not constrained by the DDH potential, which actually results from the self-conserved $\rho\rho\gamma$ vertex mentioned above, while the last equality shows the transversality of NB currents. Obviously, the total PNC EM EC operator, $(\rho_{PNC}^{(2)}, \mathbf{j}_{PNC}^{(2)})$ we construct, Eqs. (17-24), satisfies the total current conservation condition

$$[\rho^{(1)} + \rho_{PNC}^{(2)}, T + V_{PC} + V_{PNC}] = \mathbf{k} \cdot (\mathbf{j}^{(1)} + \mathbf{j}_{PC}^{(2)} + \mathbf{j}_{PNC}^{(2)}),$$

as long as the two-body PC EC, $(\rho_{PC}^{(2)}, \mathbf{j}_{PC}^{(2)})$, is conserved, i.e.,

$$[\rho^{(1)}, V_{PC}] = \mathbf{k} \cdot \mathbf{j}_{PC}^{(2)},$$

($\rho_{PC}^{(2)}$ is higher order in $1/m_N$). Therefore, at least for the PNC part, we have every conservation condition met.

For the calculation of AM, we have to use both the DDH potential and current operators in coordinate space, these expressions could be found in Appendix A.

III. DEUTERON ANAPOLE MOMENT

A. Determination of the Deuteron Wave Function

Due to the PNC NN interaction, the deuteron wave function, mainly a 3S_1 state with some fraction of 3D_1 component, could have parity admixtures in $^3\tilde{P}_1$ and $^1\tilde{P}_1$ channels. The former channel, induced by the isovector part of DDH potential, is dominated by π -exchange, while the latter one, resulting from the isoscalar interaction, is purely through the heavy-meson exchange. Therefore, we express the full parity-admixed deuteron wave function as

$$\begin{aligned} \Psi(\mathbf{r}) = & \frac{1}{\sqrt{4\pi r}} \left[\left(u(r) + \frac{S_{12}(\hat{\mathbf{r}})}{\sqrt{8}} w(r) \right) \zeta_{00} \right. \\ & \left. - i\sqrt{\frac{3}{8}} (\boldsymbol{\sigma}_1 + \boldsymbol{\sigma}_2) \cdot \hat{\mathbf{r}} \tilde{v}_{3p1}(r) \zeta_{10} + i\frac{\sqrt{3}}{2} (\boldsymbol{\sigma}_1 - \boldsymbol{\sigma}_2) \cdot \hat{\mathbf{r}} \tilde{v}_{1p1}(r) \zeta_{00} \right] \chi_{1J_z}, \end{aligned} \quad (33)$$

where $S_{12}(\hat{\mathbf{r}}) \equiv 3\boldsymbol{\sigma}_1 \cdot \hat{\mathbf{r}} \boldsymbol{\sigma}_2 \cdot \hat{\mathbf{r}} - \boldsymbol{\sigma}_1 \cdot \boldsymbol{\sigma}_2$ and χ and ζ represents spinor and isospinor respectively.

The Schrödinger equation in the center of mass frame is

$$H \Psi(\mathbf{r}) = \left[-\frac{1}{m_N} \left(\frac{1}{r} \frac{\partial^2}{\partial r^2} r - \frac{l(l+1)}{r^2} \right) + V_C(r) + V_T(r) S_{12}(\hat{\mathbf{r}}) + V_{PNC}(\mathbf{r}) \right] \Psi(\mathbf{r})$$

$$= E \Psi(\mathbf{r}), \quad (34)$$

where $V_C(r)$ and $V_T(r)$ are the central and tensor parts of the strong potential, respectively, and $V_{PNC}(\mathbf{r})$ is the sum of the PNC potentials given above. Up to the linear order of PNC NN interaction, radial wave functions of each channel satisfy the differential equations

$$u''(r) + m_N (E - V_C(r)) u(r) = \sqrt{8} m_N V_T(r) w(r), \quad (35)$$

$$w''(r) - \frac{6}{r^2} w(r) - m_N (E - V_C(r) + 2V_T(r)) w(r) = \sqrt{8} m_N V_T(r) u(r), \quad (36)$$

$$\begin{aligned} \tilde{v}_{3p1}''(r) - \frac{2}{r^2} \tilde{v}_{3p1}(r) + m_N (E - V_C(r) - 2V_T(r)) \tilde{v}_{3p1}(r) = \\ \frac{2}{\sqrt{3}} \left[\left(u(r) + \frac{1}{\sqrt{2}} w(r) \right) \frac{\partial}{\partial r} \left(F_\pi^1(r) + \sqrt{2} F_\rho^1(r) - \sqrt{2} F_\omega^1(r) \right) \right. \\ \left. + 2\sqrt{2} (F_\rho^1(r) - F_\omega^1(r)) \frac{\partial}{\partial r} \left(u(r) + \frac{1}{\sqrt{2}} w(r) \right) - \frac{2\sqrt{2}}{r} (F_\rho^1(r) - F_\omega^1(r)) (u(r) - \sqrt{2} w(r)) \right] \end{aligned} \quad (37)$$

$$\begin{aligned} \tilde{v}_{1p1}''(r) - \frac{2}{r^2} \tilde{v}_{1p1}(r) + m_N (E - V_C(r) + 4V_T(r)) \tilde{v}_{1p1}(r) = \\ \frac{2}{\sqrt{3}} \left[\left(u(r) - \sqrt{2} w(r) \right) \frac{\partial}{\partial r} (3\chi_V F_\rho^0(r) - \chi_S F_\omega^0(r)) \right. \\ \left. - 2 (3F_\rho^0(r) - F_\omega^0(r)) \frac{\partial}{\partial r} (u(r) - \sqrt{2} w(r)) + \frac{2}{r} (3F_\rho^0(r) - F_\omega^0(r)) (u(r) + 2\sqrt{2} w(r)) \right], \end{aligned} \quad (38)$$

where $F_\pi^1(r) \equiv g_{\pi NN} h_\pi^1 f_\pi(r)$, $F_\rho^0(r) \equiv g_{\rho NN} h_\rho^0 f_\rho(r)$, $F_\rho^1(r) \equiv g_{\rho NN} h_\rho^1 f_\rho(r)$, $F_\omega^0(r) \equiv g_{\omega NN} h_\omega^0 f_\omega(r)$, and $F_\omega^1(r) \equiv g_{\omega NN} h_\omega^1 f_\omega(r)$. In our numerical calculations, $g_{\pi NN} = 13.45$, $g_{\rho NN} = 2.79$, $g_{\omega NN} = 8.37$, as well as DDH best values (in units of 10^{-7}): $h_\pi^1 = 4.6$, $h_\rho^0 = -11.4$, $h_\rho^1 = -0.2$, $h_\rho^2 = -9.5$, $h_\omega^0 = -1.9$, and $h_\omega^1 = -1.1$, are assumed.

B. Anapole Moment : Expressions of Matrix Elements

The anapole operator we use takes the form

$$\mathbf{a} \equiv \frac{2\pi}{3} \int d\mathbf{x} \, \mathbf{x} \times (\mathbf{x} \times \mathbf{j}(\mathbf{x})), \quad (39)$$

where $\mathbf{j}(\mathbf{x})$ contains all the currents discussed in Sec. II. Note that this form is equivalent to what have been recommended in Refs. [9, 25], a result from implementing the extended Siegert's theorem [26].

With the deuteron wave function, Eq. (33), we obtain the anapole moment for the spin term;

$$\begin{aligned} \mathbf{a}_{spin} = -\frac{\pi}{\sqrt{6} m_N} \left[\mu_V \int dr \, r \left(u(r) - \sqrt{2} w(r) \right) \tilde{v}_{3p1}(r) \right. \\ \left. - \sqrt{2} \mu_S \int dr \, r \left(u(r) + \frac{1}{\sqrt{2}} w(r) \right) \tilde{v}_{1p1}(r) \right] e \mathbf{I}, \end{aligned} \quad (40)$$

where $\mathbf{I} \equiv \frac{1}{2}\chi_{1J_z}^\dagger(\boldsymbol{\sigma}_1 + \boldsymbol{\sigma}_2)\chi_{1J_z}$ is the intrinsic spin taken in the spinor basis, and we note that this is equivalent to the total angular momentum taken in the total angular momentum basis, i.e., $\mathbf{I} = \langle J=1, J_z | \mathbf{J} | J=1, J_z \rangle$ ².

The matrix element of the convection current is written as

$$\begin{aligned} \mathbf{j}_{conv}(\mathbf{x}) &= \mathbf{j}_{conv}^+(\mathbf{x}) + \mathbf{j}_{conv}^-(\mathbf{x}), \\ \mathbf{j}_{conv}^+(\mathbf{x}) &\equiv \frac{e}{4m_N} \int d\mathbf{r}_1 d\mathbf{r}_2 \frac{1}{\sqrt{4\pi r}} \chi_{1J_z}^\dagger \left(u(r) + \frac{S_{12}(\hat{\mathbf{r}})}{\sqrt{8}} w(r) - i\frac{\sqrt{3}}{2}(\boldsymbol{\sigma}_1 - \boldsymbol{\sigma}_2) \cdot \hat{\mathbf{r}} \tilde{v}_{1p1}(r) \right) \\ &\quad \times (\mathbf{p}_1, \mathbf{p}_2)^+ \left(u(r) + \frac{S_{12}(\hat{\mathbf{r}})}{\sqrt{8}} w(r) + i\frac{\sqrt{3}}{2}(\boldsymbol{\sigma}_1 - \boldsymbol{\sigma}_2) \cdot \hat{\mathbf{r}} \tilde{v}_{1p1}(r) \right) \frac{1}{\sqrt{4\pi r}} \chi_{1J_z}, \end{aligned} \quad (41)$$

$$\begin{aligned} \mathbf{j}_{conv}^-(\mathbf{x}) &\equiv \frac{e}{4m_N} \int d\mathbf{r}_1 d\mathbf{r}_2 \frac{1}{\sqrt{4\pi r}} \chi_{1J_z}^\dagger \\ &\quad \times \left[\left(u(r) + \frac{S_{12}(\hat{\mathbf{r}})}{\sqrt{8}} w(r) \right) (\mathbf{p}_1, \mathbf{p}_2)^- \left(-i\sqrt{\frac{3}{8}} \right) (\boldsymbol{\sigma}_1 + \boldsymbol{\sigma}_2) \cdot \hat{\mathbf{r}} \tilde{v}_{3p1}(r) \right. \\ &\quad \left. + i\sqrt{\frac{3}{8}} (\boldsymbol{\sigma}_1 + \boldsymbol{\sigma}_2) \cdot \hat{\mathbf{r}} \tilde{v}_{3p1}(r) (\mathbf{p}_1, \mathbf{p}_2)^- \left(u(r) + \frac{S_{12}(\hat{\mathbf{r}})}{\sqrt{8}} w(r) \right) \right] \frac{1}{\sqrt{4\pi r}} \chi_{1J_z}, \end{aligned} \quad (42)$$

where

$$(\mathbf{p}_1, \mathbf{p}_2)^\pm \equiv \{\mathbf{p}_1, \delta^3(\mathbf{x} - \mathbf{r}_1)\} \pm \{\mathbf{p}_2, \delta^3(\mathbf{x} - \mathbf{r}_2)\}. \quad (43)$$

As been argued in [8]

$$\frac{2\pi}{3} \int d\mathbf{x} \mathbf{x} \times (\mathbf{x} \times \mathbf{j}_{conv}^-(\mathbf{x})) = \left\langle -i\frac{\pi e}{12m_N} [\mathbf{l}^2, \mathbf{r}] \right\rangle, \quad (44)$$

where $\mathbf{l} = \mathbf{r} \times \mathbf{p}$ ³. Indeed, it can be shown that the matrix element of the operator $(\mathbf{p}_1, \mathbf{p}_2)^+$ is proportional to

$$-2i\mathbf{R} + \mathbf{R}(\mathbf{R} \cdot \mathbf{P}) + \mathbf{r} \left(\mathbf{R} \cdot \mathbf{p} + \frac{1}{4}\mathbf{r} \cdot \mathbf{P} \right) - \left(\mathbf{R}^2 + \frac{1}{4}\mathbf{r}^2 \right) \mathbf{P} - 2\mathbf{R} \cdot \mathbf{r} \mathbf{p},$$

where $\mathbf{R} = (\mathbf{r}_1 + \mathbf{r}_2)/2$ is the coordinate of the center of mass and $\mathbf{P} = \mathbf{p}_1 + \mathbf{p}_2$ is its conjugate momentum. In the center of mass frame where only the relative coordinate or momentum are relevant, the terms proportional to $\mathbf{R}$ or $\mathbf{P}$ are discarded. Consequently, the contribution from $(\mathbf{p}_1, \mathbf{p}_2)^+$ is zero and the magnitude from the convection current is determined by Eq. (42). The anapole moment contributed by the convection term then reads

$$\mathbf{a}_{conv} = \frac{1}{3} \frac{\pi}{\sqrt{6} m_N} \int dr r \left(u(r) - \sqrt{2}w(r) \right) \tilde{v}_{3p1}(r) e \mathbf{I}. \quad (45)$$

² Since the anapole moment is a vector moment, it should be proportional to the total angular momentum which is the only intrinsic vector of a system.

³ We stress that the equality holds for the matrix elements and the operators themselves do not satisfy the equality.

Contributions from the PNC exchange currents are evaluated with the parity-even channels (3S_1 and 3D_1) in the initial and final states. These channels are in spin-triplet, $S = 1$, and isospin-singlet, $T = 0$, state and for these states we have the spin and isospin selections rules

$$\langle S = 1 || (\boldsymbol{\sigma}_1 - \boldsymbol{\sigma}_2) || S = 1 \rangle = \langle S = 1 || (\boldsymbol{\sigma}_1 \times \boldsymbol{\sigma}_2) || S = 1 \rangle = 0, \quad (46)$$

$$\langle T = 0 || (\boldsymbol{\tau}_1 \times \boldsymbol{\tau}_2)^z || T = 0 \rangle = 0, \quad (47)$$

where the double bar $||$ means the reduced matrix element. It is easily seen that the matrix elements of $\mathbf{j}_{pair+KR}^{\rho^\pm}$, $\mathbf{j}_{mesonic}^{\rho^\pm}$, $\mathbf{j}_{mesonic}^{\rho\pi}$, and the part proportional to $\boldsymbol{\sigma}_1 - \boldsymbol{\sigma}_2$ in $\mathbf{j}_{pair}^{\rho^0}$ and $\mathbf{j}_{pair}^\omega$ vanish. Then we are left with π -pair, π -mesonic, and the parts proportional to $\boldsymbol{\sigma}_1 + \boldsymbol{\sigma}_2$ in ρ^0 -pair and ω -pair terms. Pair and mesonic pion contributions were already calculated in [8] which are given as

$$\mathbf{a}_{pair}^\pi = -\frac{\sqrt{2}\pi g_{\pi NN}}{9m_N} \int dr r^2 f_\pi(r) \left(u(r) + \frac{1}{\sqrt{2}}w(r) \right) \left(u(r) - \sqrt{2}w(r) \right) e \mathbf{I} h_\pi^1, \quad (48)$$

$$\begin{aligned} \mathbf{a}_{mesonic}^\pi &= \frac{\sqrt{2}\pi g_{\pi NN}}{3m_N m_\pi} \int dr r f_\pi(r) \left(u(r) + \frac{1}{\sqrt{2}}w(r) \right) \\ &\times \left[u(r) \left(1 - \frac{1}{3}m_\pi r \right) - \frac{1}{\sqrt{2}}w(r) \left(1 + \frac{1}{3}m_\pi r \right) \right] e \mathbf{I} h_\pi^1, \end{aligned} \quad (49)$$

and ρ^0 - and ω -meson contributions read

$$\mathbf{a}_{pair}^{\rho^0} = -\frac{2\pi g_{\rho NN}}{9m_N} \int dr r^2 f_\rho(r) \left(u(r) + \frac{1}{\sqrt{2}}w(r) \right) \left(u(r) - \sqrt{2}w(r) \right) e \mathbf{I} h_\rho^1, \quad (50)$$

$$\mathbf{a}_{pair}^\omega = \frac{2\pi g_{\omega NN}}{9m_N} \int dr r^2 f_\omega(r) \left(u(r) + \frac{1}{\sqrt{2}}w(r) \right) \left(u(r) - \sqrt{2}w(r) \right) e \mathbf{I} h_\omega^1. \quad (51)$$

C. Numerical Results for the Argonne v_{18} NN Interaction Model

With the Av_{18} model, we obtain the numerical results (in units of fm^2)

$$\mathbf{a}_{spin} = -0.547 h_\pi^1 e \mathbf{I} + (-3.7 h_\rho^1 + 10.8 h_\omega^1 + 7.3 h_\rho^0 + 3.7 h_\omega^0) \times 10^{-3} e \mathbf{I}, \quad (52)$$

$$\mathbf{a}_{conv} = 0.039 h_\pi^1 e \mathbf{I} + (2.7 h_\rho^1 - 7.6 h_\omega^1) \times 10^{-4} e \mathbf{I}, \quad (53)$$

$$\mathbf{a}_{ex}^\pi = \mathbf{a}_{pair}^\pi + \mathbf{a}_{mesonic}^\pi = (-0.027 + 0.028) h_\pi^1 e \mathbf{I}, \quad (54)$$

$$\mathbf{a}_{ex}^\rho = \mathbf{a}_{pair}^{\rho^0} = -0.7 \times 10^{-4} h_\rho^1 e \mathbf{I}, \quad (55)$$

$$\mathbf{a}_{ex}^\omega = \mathbf{a}_{pair}^\omega = 1.8 \times 10^{-4} h_\omega^1 e \mathbf{I}. \quad (56)$$

The contribution from the nucleonic AM, $\mathbf{a}_N$, should also be included in the full result. Since the deuteron is an isosinglet state, only the isoscalar component of the nucleonic AM contributes, i.e.,

$$\mathbf{a}_N = \langle d | \sum_{i=1}^2 (a_S^{(1)} + a_V^{(1)} \tau_i^z) \boldsymbol{\sigma}_i | d \rangle = 2 a_S^{(1)} \left(1 - \frac{3}{2} P_D \right) \mathbf{I}, \quad (57)$$

where $a_{S,V}^{(1)}$ denote the isoscalar and isovector nucleonic AM, and P_D is the probability of deuteron D -state. Several theoretical estimates for the nucleonic AM exist [25, 27, 28, 29, 30, 31], and here we use the result of Ref. [31] because it is the most recent one which includes the full DDH interaction at the nucleon level.

For the pion sector, Ref. [31] gave

$$a_S^{\pi(1)} = -\frac{g_A h_\pi^1}{12\sqrt{2}f_\pi m_\pi} \frac{\Lambda_\chi^2}{m_N^2} \left(1 - \frac{6 m_\pi}{\pi m_N} \ln \frac{\Lambda_\chi}{m_\pi}\right),$$

where Λ_χ is the chiral symmetry breaking scale (the authors chose it to be $4\pi f_\pi$)⁴. When setting $\Lambda_\chi = m_N$, the leading term is equivalent to what has been used in Refs. [4, 5, 8] for the deuteron AM calculations, while the full result is the same as been used in Ref. [7]. By including the heavy mesons, the numerical result is [31]

$$a_S^{(1)} = -0.274 h_\pi^1 - 0.419 h_\rho^1 - 0.129 h_\omega^0, \quad (58)$$

and this leads to the nucleonic contribution

$$\mathbf{a}_N = (-0.250 h_\pi^1 - 0.383 h_\rho^1 - 0.118 h_\omega^0) e \mathbf{I}. \quad (59)$$

Although we do not have the exact PC EC corresponding to Av_{18} potential, we try to approximate this by the EC coming out from one-pion exchange diagrams. Corresponding currents in the configuration space are given in the Appendix A. Pair- and mesonic-term of the PC EC contributions read

$$\mathbf{a}_{pair}^{PC} = \frac{1}{3\sqrt{6}} \left(\frac{g_{\pi NN}}{2m_N}\right)^2 \int dr e^{-m_\pi r} (1 + m_\pi r) \tilde{v}_{3p1}(r) \left(u(r) - \sqrt{2}w(r)\right) e \mathbf{I}, \quad (60)$$

$$\mathbf{a}_{mesonic}^{PC} = -\frac{1}{3\sqrt{6}} \left(\frac{g_{\pi NN}}{2m_N}\right)^2 \int dr e^{-m_\pi r} \tilde{v}_{3p1}(r) \left[3u(r) - m_\pi r \left(u(r) + \frac{1}{\sqrt{2}}w(r)\right)\right] e \mathbf{I}. \quad (61)$$

Numerically we obtain

$$\begin{aligned} \mathbf{a}_{ex}^{PC} &= \mathbf{a}_{pair}^{PC} + \mathbf{a}_{mesonic}^{PC} \\ &= -(7.8 h_\pi^1 + 0.4 h_\rho^1 - 1.1 h_\omega^1) \times 10^{-4} e \mathbf{I}. \end{aligned} \quad (62)$$

In the next section, we will discuss if this is a reasonable approximation and to what extent is the breaking of current conservation due to the non-exact PC EC.

⁴ Note that a factor of $4\pi/m_N^2$ is needed for converting the dimensionless AM defined in Lorentz-Heaviside units into our definition.




Figure 3: The left panel shows the behavior of $r^2(u(r) + w(r)/\sqrt{2})(u(r) - \sqrt{2}w(r))$. The central panel compares the Yukawa function of the pion (solid line) and rho-meson (dotted line). The right panel is the behavior of the integrand for the π^- and ρ^- pair terms in Eq. (48) and Eq. (50), respectively

Finally, the full deuteron AM could be expressed as:

$$\begin{aligned} \mathbf{a}_d &= \mathbf{a}_{spin} + \mathbf{a}_{conv} + \mathbf{a}_{ex}^{PNC} + \mathbf{a}_N + \mathbf{a}_{ex}^{PC} \\ &= (-0.756 h_\pi^1 - 0.387 h_\rho^1 + 0.010 h_\omega^1 + 0.007 h_\omega^0 - 0.114 h_\omega^0) e \mathbf{I}. \end{aligned} \quad (63)$$

IV. DISCUSSIONS

Presenting the results, we separated the contributions of the pion and the heavy mesons so that their relative order of magnitude can be compared easily. In the result of the nuclear part, i.e. the contributions from deuteron wave function or exchange currents, which include spin, convection, PNC EC and PC EC, the magnitudes of the heavy meson contributions are commonly suppressed by an order of 2 or 3. This suppression can be understood most easily from Eqs. (48, 50, 51), PNC pair terms of π , ρ and ω mesons, respectively. Aside from the weak coupling constants, they differ only by the Yukawa functions $f_\pi(r)$ or $f_{\rho,\omega}(r)$ in the integrand. At small r , the deuteron wave function is proportional to r (for the simplicity of argument, we neglect D -state wave function) and increases very fast up to $r \sim 2$ fm at which the maximum is reached. Afterwards, the wave function decreases very slowly, converging to zero. The factor r^2 multiplied by $u^2(r)$ gives rise to the suppression at short range. The left panel of Fig. 3 shows this behavior. One can expect that this r^4 behavior at short distances makes the contribution from $r \leq 1$ fm quite small. On the contrary, the Yukawa function, which behaves like $1/r$ at small r and also depends strongly on the mass of the meson, is short-range peaked. A comparison of $f_\pi(r)$ and $f_\rho(r)$ is given in the central panel of Fig. 3. The quantity $f_\rho(r)$ is non-negligible for about $r \leq 0.5$ fm but the remaining part of the integrand, approximately $r^2 u^2(r)$ is small in this region. Consequently the heavy meson will be suppressed substantially compared to π . The right panel of Fig. 3 shows the behavior

term	spin	conv.	PNC EC	PC EC	nucleonic	total
$(\rho + \omega)/\pi$	4.0	0.6	-2.0	-3.1	-26.2	-5.9

Table I: Comparison of the heavy meson with the pion contribution to the deuteron anapole moment, term by term. The ratios are given in %.

of the total integrands in Eqs. (48, 50). If we approximate the integrand as $r^3 \exp(-m_X r)$, the maximum of the integrand is reached at $r \sim 3/m_X$. This approximation is good for ρ but bad for π , however, this can provide a semi-quantitative explanation for the suppression of the heavy mesons. At the maximum, the integrand has the value proportional to $1/m_X^3$. If comparison is made for π and ρ , the maximum value of ρ is smaller than π by a factor of $1/200$, which can account for the suppression of order 2 of the heavy mesons.

To a lesser extent, a similar argument can be applied to the qualitative understanding of the heavy-meson suppression of spin and convection terms. The right hand side of Eqs. (37, 38) which are the sources of parity admixed P -states also contains the Yukawa functions. When the equations are solved, i.e. integrated with respect to r , one can expect some amount of suppression for the heavy mesons as was discussed in the analysis of pair EC's, but without the factor r^2 . In Tab. I, we summarize the ratio of the heavy meson contribution to the pion with the weak coupling constants given by DDH best values. The magnitudes of the heavy meson terms are suppressed by an order of two compared to the pion terms in common. As far as the nuclear part is concerned, π contribution is by far dominant. If one can disentangle the nuclear part contribution from the total deuteron anapole moment, this can provide information to extract the magnitude of h_π^1 with high accuracy. As a side remark, it is noticed that the combination of the isovector couplings, h_π^1 , h_ρ^1 and h_ω^1 , entering the spin and convection contributions is close to the one determining low energy PNC nucleon-nucleon scattering, roughly $3 h_\pi^1 + 0.02 h_\rho^1 - 0.06 h_\omega^1$ [32]. This lets us think that the Danilov's approach [33], whose application for the deuteron anapole moment was proposed later on by Savage [6], should provide a reasonable estimate for this contribution.

Contrary to the nuclear part, the nucleonic one has sizable contribution from the heavy mesons. The relative ratio of the heavy meson is about 26% of the pion based on the nucleonic anapole estimate of Ref. [31]. However, we should also note that these authors considered more non-DDH type couplings such as non-Yukawa type πNN couplings and the inclusion of hyperons. So far, no detailed knowledge about these exotic couplings exist, and the theoretical uncertainty could be huge. The study of these terms and their implication for two-body nuclear contributions will be an

interesting topic for further exploration.

Another issue that should be addressed is the gauge invariance of the results. We showed in Sect. II that the PNC ECs we constructed satisfy the gauge invariance with the DDH potential. However, with the phenomenological strong interaction models like the one adopted in this work, gauge invariance may not be satisfied if the ECs are not consistent with the phenomenological potentials. For Av_{18} , the potential contains 18 types of operators but we took into account the dominant ECs only, which naturally leads to inconsistency, or breakdown of gauge invariance. Even though the inconsistency may only change the result slightly, the investigation of the extent to which gauge invariance is broken could help to get an estimate of the error.

Most phenomenological potentials are very complicated and the analytic analysis of gauge invariance is a formidable or practically impossible work. However as suggested in [8], one can estimate the amount of gauge invariance breaking by comparing the anapole moment obtained from Eq. (39) and

$$\mathbf{a} = -\pi \int d\mathbf{x} x^2 \mathbf{j}(\mathbf{x}), \quad (64)$$

which can be obtained from Eq. (39) with the assumption

$$\nabla \cdot \mathbf{j}(\mathbf{x}) = 0. \quad (65)$$

In [8], we observed that the inclusion of the PC ECs of π removes almost all the inconsistency. In this work, we follow the same procedure to investigate the breakdown of gauge invariance.

The spin current can be shown easily to satisfy Eq. (65). The convection, pair and mesonic anapole moment of π , ρ - and ω -mesons with the definition given by Eq. (64) can be calculated straightforwardly to give

$$\mathbf{a}_{conv}^{CC} = \frac{\pi}{2\sqrt{6}m_N} \int dr r^2 \left[u(r) \left(\tilde{v}'_{3p1}(r) + 2\frac{\tilde{v}_{3p1}(r)}{r} \right) + \frac{w(r)}{\sqrt{2}} \left(\tilde{v}'_{3p1}(r) - \frac{\tilde{v}_{3p1}(r)}{r} \right) \right] e \mathbf{I}, \quad (66)$$

$$\mathbf{a}_{pair}^{\pi,CC} = -\frac{\pi g_{\pi NN}}{2\sqrt{2}m_N} \int dr r^2 f_{\pi}(r) \left(u^2(r) - \frac{w^2(r)}{2} \right) h_{\pi}^1 e \mathbf{I}, \quad (67)$$

$$\begin{aligned} \mathbf{a}_{mesonic}^{\pi,CC} &= \frac{g_{\pi NN}}{24\sqrt{2}m_N m_{\pi}} \int dr e^{-m_{\pi}r} \\ &\times \left[\left(u^2(r) - \frac{w^2(r)}{2} \right) (m_{\pi}r + 4) - \frac{1}{3} \left(u(r) + \frac{w(r)}{\sqrt{2}} \right)^2 m_{\pi}r (m_{\pi}r + 3) \right] h_{\pi}^1 e \mathbf{I}, \quad (68) \end{aligned}$$

$$\mathbf{a}_{pair}^{\rho^0,CC} = -\frac{\pi g_{\rho NN}}{2m_N} \int dr r^2 f_{\rho}(r) \left(u^2(r) - \frac{w^2(r)}{2} \right) h_{\rho}^1 e \mathbf{I}, \quad (69)$$

$$\mathbf{a}_{pair}^{\omega,CC} = \frac{\pi g_{\omega NN}}{2m_N} \int dr r^2 f_{\omega}(r) \left(u^2(r) - \frac{w^2(r)}{2} \right) h_{\omega}^1 e \mathbf{I}, \quad (70)$$

	Eq. (39)			Eq. (64)		
	h_π^1	h_ρ^1	h_ω^1	h_π^1	h_ρ^1	h_ω^1
conv.	0.03925	0.00027	-0.00076	0.05348	0.00024	-0.00066
PNC π pair	-0.02668			-0.07895		
PNC π mesonic	0.02830			0.04706		
PNC ρ pair		-0.00007			-0.00023	
PNC ω pair			0.00018			0.00061
PC π pair	0.01400	0.00013	-0.00038	0.06325	0.00062	-0.00177
PC π mesonic	-0.01478	-0.00017	0.00049	-0.04366	-0.00047	0.00133
Total-PC	0.04087	0.00020	-0.00058	0.02159	0.00001	-0.00005
Total	0.04010	0.00016	-0.00047	0.04117	0.00017	-0.00049

Table II: Coefficients of the weak coupling constants for given terms with different definitions of the anapole operator, Eq. (39) and Eq. (64).

$$\mathbf{a}_{pair}^{PC,CC} = \frac{1}{\sqrt{6}} \left(\frac{g_{\pi NN}}{2m_N} \right)^2 \int dr e^{-m_\pi r} (1 + m_\pi r) \tilde{v}_{3p1}(r) \left(u(r) - \frac{1}{\sqrt{8}} w(r) \right) e \mathbf{I}, \quad (71)$$

$$\begin{aligned} \mathbf{a}_{mesonic}^{PC,CC} = & -\frac{1}{12\sqrt{6}} \left(\frac{g_{\pi NN}}{2m_N} \right)^2 \int dr e^{-m_\pi r} \tilde{v}_{3p1}(r) \\ & \times \left[u(r)(18 + 2m_\pi r - m_\pi^2 r^2) - \frac{w(r)}{\sqrt{2}} m_\pi r (4 + m_\pi r) \right] e \mathbf{I}, \end{aligned} \quad (72)$$

where the superscript CC denotes the quantity calculated with Eq. (64). Numerical results are summarized in Tab. II.

Judging from the "Total" column in Table II, two definitions of the anapole operator only differ in results by 3%, 6%, and 4% for the π -, ρ -, and ω -exchange respectively; this means our result satisfy current conservation very well. This is quite surprising because only the PC one-pion exchange current—not fully consistent with the adopted strong potential, Av_{18} —is included in our calculation. The reason is partly because the AM is a r^2 -weighted moment, plus the deuteron wave function peaks around 2 fm with a long tail; therefore the long-range physics, which is dominated by the one-pion exchange including the case of Av_{18} , becomes much more important. This explains why our conclusion differs from the one found in Ref. [13] where large effects due to undetermined ECs were found. Their observation was concerning a different operator (E1 transition), which has a range relatively shorter than the quantity considered in this work.

Furthermore, one can observe that for the contributions from exchange currents, Eq. (39) always give smaller values than Eq. (64). This implies the calculation using Eq. (39) suffers less from

the incomplete knowledge or uncertainty of exchange currents which causes the breaking of current conservation. For example, if the PC pion ECs are left out in our calculation, by comparing the "Total" and "Total-PC" columns in Table II, the error is 2%, 25%, and 23% for π , ρ , and ω component respectively when Eq. (39) is used; while the error is -46%, -94%, and -90% for Eq. (64). Perhaps, the reason for the small contribution of ECs in the former case is to be looked for in the proportionality of the dominant ECs to the position vector $\vec{x}$, which readily gives zero when inserted into the corresponding anapole operator, Eq. (39). Therefore the merit of Eq. (39) is justified, at least for the case of deuteron.

In conclusion, we constructed the PNC ECs due to one π -, ρ - and ω -exchange, and showed that they satisfy the current conservation; and hence are fully consistent with the adopted DDH PNC potential. An application was made to the calculation of the deuteron anapole moment. We observed that, for the nuclear part, the contribution of heavy mesons is suppressed by an order of two compared to the pion, a result consistent with the similar work by Blunden [34]. Consistency with Av_{18} was also checked. We found that the approximation of using only the PC one-pion exchange current is pretty good—the breaking of current conservation only amounts to a few percent, and this should be fixed by using the PC exchange current fully consistent with Av_{18} . Therefore, the contribution from the nuclear part to the deuteron anapole moment can be determined with an error less than 5%, while the major uncertainty should come from the nucleonic anapole moment instead.

Acknowledgements

Work of C.H.H. is partially supported by KOSEF (Grant No. 2000-2-11100-004-4).

Appendix A: PNC NN POTENTIAL AND EXCHANGE CURRENTS IN COORDINATE SPACE

The DDH potential in coordinate space could easily be obtained by applying the following transformation rules to Eqs. (25-27)

$$\begin{aligned}\mathbf{u}_X &\rightarrow \mathbf{u}_X(\mathbf{r}) = [\mathbf{p}, f_X(r)], \\ \mathbf{v}_X &\rightarrow \mathbf{v}_X(\mathbf{r}) = \{\mathbf{p}, f_X(r)\},\end{aligned}$$

where $\mathbf{r} \equiv \mathbf{r}_1 - \mathbf{r}_2$; $r = |\mathbf{r}|$; $\mathbf{p} \equiv (\mathbf{p}_1 - \mathbf{p}_2)/2 = -i\nabla_{\mathbf{r}}$; and the Yukawa functions $f_X(r)$ are defined as

$$f_X(r) = \frac{e^{-m_X r}}{4\pi r}.$$

For the PNC MECs, we list all the leading-order, $O(1/m_N)$, 3-currents, which are relevant for the AM calculation:

$$\mathbf{j}_{pair}^\pi(\mathbf{x}; \mathbf{r}_1, \mathbf{r}_2) = -\frac{e g_{\pi NN} h_\pi^1}{2\sqrt{2} m_N} (\boldsymbol{\tau}_1 \cdot \boldsymbol{\tau}_2 - \tau_1^z \tau_2^z) f_\pi(r) \sum_{i=1}^2 \delta^{(3)}(\mathbf{x} - \mathbf{r}_i) \boldsymbol{\sigma}_i, \quad (\text{A1})$$

$$\begin{aligned} \mathbf{j}_{mesonic}^\pi(\mathbf{x}; \mathbf{r}_1, \mathbf{r}_2) = & -\frac{e g_{\pi NN} h_\pi^1}{2\sqrt{2} m_N} (\boldsymbol{\tau}_1 \cdot \boldsymbol{\tau}_2 - \tau_1^z \tau_2^z) \\ & \times (\nabla_1 - \nabla_2) [(\boldsymbol{\sigma}_1 \cdot \nabla_1 - \boldsymbol{\sigma}_2 \cdot \nabla_2), f_\pi(r_{x1}) f_\pi(r_{x2})], \end{aligned} \quad (\text{A2})$$

$$\begin{aligned} \mathbf{j}_{pair+KR}^{\rho^\pm}(\mathbf{x}; \mathbf{r}_1, \mathbf{r}_2) = & -\frac{e g_{\rho NN}}{2 m_N} f_\rho(r) \left(h_\rho^0 - \frac{1}{2\sqrt{6}} h_\rho^2 \right) \\ & \times \left((\boldsymbol{\tau}_1 \cdot \boldsymbol{\tau}_2 - \tau_1^z \tau_2^z) (\boldsymbol{\sigma}_1 - \boldsymbol{\sigma}_2) \left(\delta^{(3)}(\mathbf{x} - \mathbf{r}_1) - \delta^{(3)}(\mathbf{x} - \mathbf{r}_2) \right) \right. \\ & \left. + (1 + \chi_V) (\boldsymbol{\tau}_1 \times \boldsymbol{\tau}_2)^z (\boldsymbol{\sigma}_1 \times \boldsymbol{\sigma}_2) \sum_i \delta^{(3)}(\mathbf{x} - \mathbf{r}_i) \right), \end{aligned} \quad (\text{A3})$$

$$\begin{aligned} \mathbf{j}_{pair}^{\rho^0}(\mathbf{x}; \mathbf{r}_1, \mathbf{r}_2) = & -\frac{e g_{\rho NN}}{2 m_N} f_\rho(r) \tau_1^z \tau_2^z \\ & \times \left(\left(h_\rho^0 + \frac{1}{2} h_\rho^1 (\tau_1^z + \tau_2^z) + \frac{1}{\sqrt{6}} h_\rho^2 \right) (\boldsymbol{\sigma}_1 - \boldsymbol{\sigma}_2) + \frac{1}{2} h_\rho^1 (\tau_1^z - \tau_2^z) (\boldsymbol{\sigma}_1 + \boldsymbol{\sigma}_2) \right) \\ & \times \left((1 + \tau_1^z) \delta^{(3)}(\mathbf{x} - \mathbf{r}_1) - (1 + \tau_2^z) \delta^{(3)}(\mathbf{x} - \mathbf{r}_2) \right), \end{aligned} \quad (\text{A4})$$

$$\begin{aligned} \mathbf{j}_{mesonic}^{\rho^\pm}(\mathbf{x}; \mathbf{r}_1, \mathbf{r}_2) = & -\frac{e g_{\rho NN}}{2 m_N} \left(h_\rho^0 - \frac{1}{2\sqrt{6}} h_\rho^2 \right) (\boldsymbol{\tau}_1 \times \boldsymbol{\tau}_2)^z \\ & \times \left\{ (\nabla_1 - \nabla_2) \left(-i (\boldsymbol{\sigma}_1 - \boldsymbol{\sigma}_2) \cdot \{(\nabla_1 - \nabla_2), f_\rho(r_{x1}) f_\rho(r_{x2})\} \right. \right. \\ & \left. \left. + (1 + \chi_V) (\boldsymbol{\sigma}_1 \times \boldsymbol{\sigma}_2) \cdot [(\nabla_1 - \nabla_2), f_\rho(r_{x1}) f_\rho(r_{x2})] \right) \right. \\ & + 2 \nabla_x^a \left(i \{ \nabla_1^a \boldsymbol{\sigma}_2 - \nabla_2^a \boldsymbol{\sigma}_1 + \sigma_1^a \nabla_2 - \sigma_2^a \nabla_1, f_\rho(r_{x1}) f_\rho(r_{x2}) \} \right. \\ & \left. - (1 + \chi_V) [(\boldsymbol{\sigma}_1 \times \nabla_1)^a \boldsymbol{\sigma}_2 - (\boldsymbol{\sigma}_2 \times \nabla_2)^a \boldsymbol{\sigma}_1 \right. \\ & \left. \left. + \sigma_1^a \boldsymbol{\sigma}_2 \times \nabla_2 - \sigma_2^a \boldsymbol{\sigma}_1 \times \nabla_1, f_\rho(r_{x1}) f_\rho(r_{x2}) \right] \right) \\ & \left. + 4 i m_N (\boldsymbol{\sigma}_1 + \boldsymbol{\sigma}_2) [H, f_\rho(r_{x1}) f_\rho(r_{x2})] \right\} \end{aligned} \quad (\text{A5})$$

$$\mathbf{j}_{pair}^\omega(\mathbf{x}; \mathbf{r}_1, \mathbf{r}_2) = -\frac{e g_{\omega NN}}{2 m_N} f_\omega(r)$$

$$\times \left(\left(h_\omega^0 + \frac{1}{2} h_\omega^1 (\tau_1^z + \tau_2^z) \right) (\boldsymbol{\sigma}_1 - \boldsymbol{\sigma}_2) + \frac{1}{2} h_\omega^1 (\tau_1^z - \tau_2^z) (\boldsymbol{\sigma}_1 + \boldsymbol{\sigma}_2) \right) \\ \times \left((1 + \tau_1^z) \delta^{(3)}(\mathbf{x} - \mathbf{r}_1) - (1 + \tau_2^z) \delta^{(3)}(\mathbf{x} - \mathbf{r}_2) \right), \quad (\text{A6})$$

$$\mathbf{j}_{mesonic}^{\rho\pi}(\mathbf{x}; \mathbf{r}_1, \mathbf{r}_2) = -\frac{e g_{\rho NN} g_{\rho\pi\gamma} h_\pi^1}{\sqrt{2} m_\rho} (\boldsymbol{\tau}_1 \times \boldsymbol{\tau}_2)^z \\ \times (\nabla_1 \times \nabla_2) \left(f_\rho(r_{x1}) f_\pi(r_{x2}) + f_\pi(r_{x1}) f_\rho(r_{x2}) \right), \quad (\text{A7})$$

where $r_{xi} \equiv |\mathbf{x} - \mathbf{r}_i|$; ∇_i and ∇_x act on the source point $\mathbf{r}_i$ and field point $\mathbf{x}$ respectively; and the superscript a is the index to be summed from 1 to 3. Note that we separate charged and neutral ρ mesons according to their isospin structure; and the last term of Eq. (A5) should be combined with a charge density corresponding to Eq. (24) in order to insure the current conservation.

PC one-pion ECs of the pair and mesonic terms that contribute to the AM read

$$\mathbf{j}_{pair}^{PC}(\mathbf{x}; \mathbf{r}_1, \mathbf{r}_2) = \left(\frac{g_{\pi NN}}{2m_N} \right)^2 (\boldsymbol{\tau}_1 \times \boldsymbol{\tau}_2)^z \\ \times \left[\boldsymbol{\sigma}_2 \delta^{(3)}(\mathbf{x} - \mathbf{r}_2) \boldsymbol{\sigma}_1 \cdot \nabla + \boldsymbol{\sigma}_1 \delta^{(3)}(\mathbf{x} - \mathbf{r}_1) \boldsymbol{\sigma}_2 \cdot \nabla \right] f_\pi(r), \quad (\text{A8}) \\ \mathbf{j}_{mesonic}^{PC}(\mathbf{x}; \mathbf{r}_1, \mathbf{r}_2) = -\left(\frac{g_{\pi NN}}{2m_N} \right)^2 (\boldsymbol{\tau}_1 \times \boldsymbol{\tau}_2)^z (\boldsymbol{\sigma}_1 \cdot \nabla_1) (\boldsymbol{\sigma}_2 \cdot \nabla_2) (\nabla_1 - \nabla_2) f_\pi(r_{x1}) f_\pi(r_{x2}). \quad (\text{A9})$$

-
- [1] Y. B. Zel'dovich, Zh. Eksp. Teor. Fiz. **33**, 1531 (1957).
 - [2] V. V. Flambaum and I. B. Khriplovich, Zh. Eksp. Teor. Fiz. **79**, 1656 (1980).
 - [3] C. S. Wood *et al.*, Science **275**, 1759 (1997).
 - [4] M. J. Savage and R. P. Springer, Nucl. Phys. A **644**, 235 (1998), erratum: *ibid.* **A 657**, 457 (1999).
 - [5] M. J. Savage and R. P. Springer, Nucl. Phys. A **686**, 413 (2001).
 - [6] M. J. Savage, Nucl. Phys. A **695**, 365 (2001).
 - [7] I. B. Khriplovich and R. V. Korkin, Nucl. Phys. A **665**, 365 (2000).
 - [8] C. H. Hyun and B. Desplanques, Phys. Lett. B **552**, 41 (2003).
 - [9] W. C. Haxton, C.-P. Liu, and M. J. Ramsey-Musolf, Phys. Rev. C **65**, 045502 (2002).
 - [10] B. Desplanques, J. F. Donoghue, and B. R. Holstein, Ann. Phys. (N.Y.) **124**, 449 (1980).
 - [11] R. B. Wiringa, V. G. J. Stoks, and R. Schiavilla, Phys. Rev. C **51**, 38 (1995).
 - [12] D. O. Riska, Phys. Rep. **181**, 207 (1989).
 - [13] R. Schiavilla, J. Carlson, and M. Paris, Phys. Rev. C **67**, 032501(R) (2003).
 - [14] M. Chemtob and M. Rho, Nucl. Phys. A **163**, 1 (1971).
 - [15] M. Chemtob, in *Mesons in Nuclei*, edited by D. Wilkinson and M. Rho (North-Holland, Amsterdam, 1979), vol. II, p. 495.

- [16] I. S. Towner, Phys. Rep. **155**, 263 (1987).
- [17] N. M. Kroll and M. A. Ruderman, Phys. Rev. **93**, 233 (1954).
- [18] B. D. Serot and J. D. Walecka, Adv. Nucl. Phys. **16**, 1 (1986).
- [19] H. Hyuga, A. Arima, and K. Shimizu, Nucl. Phys. A **336**, 363 (1980).
- [20] R. G. Arnold, C. Carlson, and F. Gross, Phys. Rev. C **21**, 1426 (1980).
- [21] S. J. Brodsky and J. R. Hiller, Phys. Rev. D **46**, 2141 (1992).
- [22] J. Adam and E. Truhlik, Czech, J. Phys. B **34**, 1157 (1984).
- [23] J.-F. Mathiot, Phys. Rep. **173**, 63 (1989).
- [24] R. J. Adler, Phys. Rev. **141**, 1499 (1966).
- [25] W. Haxton, E. M. Henley, and M. J. Musolf, Phys. Rev. Lett. **63**, 949 (1989).
- [26] J. L. Friar and S. Fallieros, Phys. Rev. C (1984).
- [27] M. J. Musolf and B. R. Holstein, Phys. Lett. B **242**, 461 (1990).
- [28] M. J. Musolf and B. R. Holstein, Phys. Rev. D **43**, 2956 (1991).
- [29] D. B. Kaplan and M. J. Savage, Nucl. Phys. A **556**, 653 (1993).
- [30] C. M. Maekawa and U. van Kolck, Phys. Lett. B **478**, 73 (2000).
- [31] S. L. Zhu, S. J. Puglia, B. R. Holstein, and M. J. Ramsey-Musolf, Phys. Rev. D **62**, 033008 (2000).
- [32] B. Desplanques, Phys. Rep. **297**, 1 (1998).
- [33] G. S. Danilov, Phys. Lett. **18**, 40 (1965).
- [34] P. G. Blunden, http://mocha.phys.washington.edu/~int_talk/WorkShops/int_02_3/People/Blunden_P/.

Note that although the full DDH potential is used, the heavy meson exchange currents are not considered in this work; and the Bonn potential is adopted as the strong interaction.